

Almost Shift Invariant Projections in Infinite Tensor Products

Ola Bratteli

Department of Mathematics, University of Oslo,
P. O. Box 1053 Blindern,
N-0316 Oslo, Norway

David E. Evans

Department of Mathematics and Computer Science,
University College of Swansea,
Singleton Park, Swansea SA2 8PP,
Wales, U.K.

Akitaka Kishimoto

Department of Mathematics,
Hokkaido University,
Sapporo, 060 Japan

The Rohlin property

Let σ be an automorphism of a unital C^* -algebra A . Following [Con], [HO], [BEK] we will say that the dynamical system (A, σ) has the *Rohlin property* if there exists a strictly increasing sequence $(n_k)_k$ of positive integers such that for each $m = n_k$ there is a sequence

$$(e_i^{(j)} | i = 0, 1, \dots, m)_j$$

of families of projections in A with

$$\begin{aligned} e_m^{(j)} &= e_0^{(j)}, \\ \sum_{i=0}^{m-1} e_i^{(j)} &= 1, \\ \lim_{j \rightarrow \infty} \|\sigma(e_i^{(j)}) - e_{i+1}^{(j)}\| &= 0 \end{aligned}$$

for $i = 0, 1, \dots, m-1$, and

$$\lim_{j \rightarrow \infty} \|[x, e_i^{(j)}]\| = 0$$

for all $x \in A$, $i = 0, 1, \dots, m-1$.

It was established in [Voi, Lemma 3.3] (see also [BKRS], Lemma 2.1) that if A is a UHF algebra (or more generally, if A is a unital AF algebra and σ is an approximately inner automorphism) and (A, σ) has the Rohlin property, then σ is an almost inductive limit automorphism. This means

that for any $B \in \mathcal{F}(A)$ = the set of finite dimensional *-subalgebras of A containing the unit of A , and any $\varepsilon > 0$, there is a $C \in \mathcal{F}(A)$ such that $B \subseteq^\varepsilon C$ and $d(\alpha(C), C) < \varepsilon$. Here we use the terminology $C_1 \subseteq^\varepsilon C_2$ iff

$$\sup\{\inf\{\|x - y\| \mid y \in C_2, \|y\| \leq 1\} \mid x \in C_1, \|x\| \leq 1\} < \varepsilon$$

and

$$d(C_1, C_2) = \inf\{\varepsilon > 0 \mid C_1 \subseteq^\varepsilon C_2 \text{ and } C_2 \subseteq^\varepsilon C_1\}$$

whenever C_1, C_2 are subspaces of A . If σ is an almost inductive limit automorphism and $\varepsilon > 0$, then there exists a unitary $v \in A$ such that $Adv \circ \sigma$ is an inductive limit automorphism and $\|v - 1\| < \varepsilon$, [Voi, Proposition 2.3]. That $\beta = Adv \circ \sigma$ is an inductive limit automorphism means that there is an increasing sequence $(A_n) \subseteq \mathcal{F}(A)$ such that $\bigcup_n A_n$ is dense in A and $\beta(A_n) = A_n$ for all n .

It is an open problem whether or not any automorphism of a UHF algebra is an almost inductive limit automorphism. If in particular $A = M_{2^\infty} = \bigotimes_{n=1}^\infty M_2$, i.e. A is the CAR algebra, and σ is the (Bernoulli) shift on A , it was established in [BKRS] that σ has the Rohlin property, and hence σ is an almost inductive limit automorphism. Let σ be a quasifree automorphism of A , i.e.

$$\sigma(a(f)) = a(Uf)$$

where $f \in \mathcal{H}$ = the one particle space, $a(f)$ is the annihilator corresponding to f , and U is a unitary operator on \mathcal{H} . It was established in [BEK] that σ has the Rohlin property if and only if $U^n - 1$ is not of Hilbert-Schmidt class for all $n \in \mathbb{Z} \setminus \{0\}$, i.e. if and only if σ^n is outer in the trace representation for all $n \in \mathbb{Z} \setminus \{0\}$. Thus σ is an almost inductive limit automorphism in this case. If on the other hand $U^n - 1$ is a Hilbert-Schmidt operator for some $n \neq 0$, then U has purely discrete spectrum and σ is a product type automorphism, and hence σ is a (strict) inductive limit automorphism.

The present paper came out of an attempt to establish that the shift σ on $A = M_{n^\infty} = \bigotimes_{n=1}^\infty M_n$ has the Rohlin property also for $n = 3, 5, \dots$ (when n is even, the Rohlin property follows from the Rohlin property of the shift on M_{2^∞}). We have, however, only been able to establish the following approximate version of the Rohlin property.

Theorem 1. *Let σ be the one-sided shift on M_{2^∞} , and let τ be the trace state on M_{n^∞} . For any finite subset $\{x_1, \dots, x_k\} \subseteq M_{n^\infty}$, any $m \in \mathbb{N}$ and any $\varepsilon > 0$ there exists projections e_0, e_1, \dots, e_{2^m} in M_{n^∞} such that*

$$\begin{aligned} e_{2^m} &= e_0, \\ e &= \sum_{i=0}^{2^m-1} e_i \text{ is a projection,} \end{aligned}$$

$$\begin{aligned}\tau(e) &> 1 - \varepsilon, \\ \|\sigma(e_i) - e_{i+1}\| &< \varepsilon\end{aligned}$$

for $i = 0, 1, \dots, 2^m - 1$, and

$$\|[x_j, e_i]\| < \varepsilon$$

for $i = 0, 1, \dots, 2^m - 1$, $j = 1, \dots, k$.

Combining Theorem 1 with the techniques of [Voi], [BKRS] we then establish that the shift is approximately an almost inductive limit automorphism in the following sense:

Corollary 2. *Let σ be the one-sided shift on M_{n^∞} . For any finite dimensional sub- $*$ -algebra D of M_{n^∞} and any $\varepsilon > 0$, there exists a projection $e \in D' \cap M_{n^\infty}$ such that*

$$\|\sigma(e) - e\| < \varepsilon$$

and

$$\tau(e) > 1 - \varepsilon,$$

and there exists a finite-dimensional $*$ -subalgebra $E \subseteq eM_{n^\infty}e$ such that

$$De \subseteq^\varepsilon E$$

and

$$d(E, \sigma(E)) < \varepsilon.$$

The embedding of $GICAR(\mathcal{H})$ into M_{n^∞}

In this section we prove Theorem 1 and Corollary 2. Let σ be the one-sided shift on M_{n^∞} , $n = 2, 3, 4, \dots$. Let $GICAR(\mathcal{H})$ be the gauge-invariant part of the CAR algebra $CAR(\mathcal{H})$ over the Hilbert space \mathcal{H} , i.e., $GICAR(\mathcal{H})$ consists of the elements in $CAR(\mathcal{H})$ which are invariant under the quasi-free action of the circle group T given by

$$a(f) \rightarrow a(zf)$$

for $z \in \mathbf{T} \subseteq \mathbf{C}$, $f \in \mathcal{H}$. Following [CE] (see also [E]), $GICAR(\mathcal{H})$ can be embedded into M_{n^∞} as follows: Let $(f_{ij})_{i,j=1}^n$ be a complete set of matrix units for M_n , and define projections e_1, e_2, \dots in $M_{n^\infty} = \bigotimes_{n=1}^\infty M_n$ as follows

$$\begin{aligned}e_1 &= \sum_{ij} \frac{1}{n} f_{ij} \otimes 1 \otimes 1 \otimes \dots, \\ e_2 &= \sum_i f_{ii} \otimes f_{ii} \otimes 1 \otimes \dots,\end{aligned}$$

$$\begin{aligned}
e_3 &= \sum_{ij} \frac{1}{n} 1 \otimes f_{ij} \otimes 1 \otimes \cdots, \\
e_4 &= \sum_i 1 \otimes f_{ii} \otimes f_{ii} \otimes \cdots,
\end{aligned}$$

etc. The e_i 's satisfy the Temperley-Lieb relations

$$\begin{aligned}
e_i e_j &= e_j e_i \quad \text{if } |i - j| \geq 2, \\
e_k e_{k \pm 1} e_k &= \frac{1}{n} e_k.
\end{aligned}$$

Furthermore,

$$\sigma(e_k) = e_{k+2}$$

for $k = 1, 2, \dots$

Let f_1, f_2, \dots be a basis for \mathcal{H} . The one-sided quasi-free shift is the injective morphism of $CAR(\mathcal{H})$ defined by

$$\beta(a_k) = a_{k+1}$$

where $a_k = a(f_k)$, $k = 1, 2, \dots$. The shift β restricts to an injective morphism of $GICAR(\mathcal{H})$ which we also denote by β .

The embedding γ of $GICAR(\mathcal{H})$ into M_{n^∞} is given by

$$\begin{aligned}
\gamma(a_i^* a_i) &= e_{2i} \\
\gamma(a_i^* a_{i+1}) &= e_{2i} (1 - n e_{2i+1}) e_{2i+2},
\end{aligned}$$

[CE]. We outline the argument from [CE]: First note that if $gl(\mathbb{N})$ denotes the Lie algebra of complex $\infty \times \infty$ matrices such that only finitely many matrix elements are non-zero, then

$$H = [H_{ij}] \in gl(\mathbb{N}) \rightarrow Q(H) = \sum_{ij} H_{ij} a_i^* a_j$$

is a Lie algebra morphism, i.e.

$$Q([H, H']) = [Q(H), Q(H')].$$

Since

$$[e_{i,i+1}, e_{i+1,i+2}] = e_{i,i+2}$$

etc., it follows that

$$[a_i^* a_{i+1}, a_{i+1}^* a_{i+2}] = a_i^* a_{i+2}$$

etc. It follows that the $*$ -algebra generated by $a_i^* a_i, a_i^* a_{i+1}$ contains all elements of the form $a_i^* a_j$, $i, j = 1, 2, \dots$. Since any gauge invariant polynomial in a_i, a_j^* , $i, j = 1, \dots$ may be written as a polynomial in $a_i^* a_j$ it follows that

We are now ready to construct the approximate Rohlin towers of Theorem 2. For economy of notation we identify $GICAR(\mathcal{H})$ with its image in M_{n_∞} .

If $\lambda, \mu \in \mathbb{T} \subseteq \mathbb{C}$, we may choose unit vectors $f, g \in \mathcal{H}$ such that f, g are orthogonal, and

$$Uf \approx \lambda f, \quad Ug \approx \mu g$$

where U is the isometry implementing the one-sided shift on \mathcal{H} . Then

$$v = a^*(f)a(g)$$

is a partial isometry with

$$\sigma(v) \approx \lambda \bar{\mu} v.$$

Furthermore

$$\begin{aligned} v^*v &= a(g)^*a(f)a(f)^*a(g) \\ &= a(f)a(f)^*a(g)^*a(g). \end{aligned}$$

Here $a(f)a(f)^*$ and $a(g)^*a(g)$ are commuting projections in different tensor factors (if f, g are chosen suitably) of trace $\frac{n-1}{n}$ and $\frac{1}{n}$, respectively, so

$$\tau(v^*v) = \frac{n-1}{n^2}$$

Also, as

$$\begin{aligned} v^*v &= a(f)^*a(g)a(g)^*a(f) \\ &= a(f)^*a(f)a(g)a(g)^*. \end{aligned}$$

and the projections $a(f)^*a(f)$, $a(g)a(g)^*$ are orthogonal to $a(f)a(f)^*$, $a(g)^*a(g)$, respectively, the projection vv^* is orthogonal to v^*v ,

$$v^*vvv^* = 0 = vv^*v^*v$$

Hence the C^* -algebra $C^*(v)$ generated by v is isomorphic to M_2 , and if $1_v = v^*v + vv^*$ is the identity of this C^* -algebra, then

$$\tau(1_v) = \frac{2(n-1)}{n^2}$$

Also

$$\sigma|_{C^*(v)} \approx \text{Ad} \begin{pmatrix} 1 & 0 \\ 0 & \lambda \bar{\mu} \end{pmatrix}$$

Now, repeat the construction with $f_2, g_2 \perp f, g$ and (f_2, g_2) orthonormal, and cut down by $1 - 1_v = 1 - vv^* - v^*v$ to obtain a partial isometry v_2 with

$$\tau(v_2v_2^*) = \tau(v_2^*v_2) = \frac{n-1}{n^2} \left(1 - 2 \frac{(n-1)}{n^2} \right)$$

and

$$\sigma(v_2) \approx \lambda \bar{\mu} v_2$$

so

$$\tau(1 - vv^* - v^*v - v_2v_2^* - v_2^*v_2) = \left(1 - 2\frac{(n-1)}{n^2}\right)^2$$

Continuing in this manner, we may construct partial isometries $v_1 = v, \dots, v_m$ such that the projections $v_1v_1^*, v_1^*v_1, v_2v_2^*, v_2^*v_2, \dots, v_mv_m^*, v_m^*v_m$ are all orthogonal, and

$$\sigma(v_k) \approx \lambda \bar{\mu} v_k$$

for all $k = 1, \dots, m$ and

$$\tau\left(1 - \sum_{k=1}^m (v_kv_k^* + v_k^*v_k)\right) = \left(1 - 2\frac{n-1}{n^2}\right)^m$$

Now, put

$$u_m = v_1 + v_2 + \dots + v_m$$

then u_m is a partial isometry such that $u_mu_m^* \perp u_m^*u_m$, $\sigma(u_m) \approx \lambda \bar{\mu} u_m$ and

$$\tau(1 - u_mu_m^* - u_m^*u_m) = \left(1 - 2\frac{n-1}{n^2}\right)^m$$

Now repeating this construction, going further and further out in $\mathcal{H} \simeq l^2(\mathbb{N})$ to find f, g 's and using $\frac{1}{2\pi i} \log(\lambda \bar{\mu}) = 2^{-1}, 2^{-2}, 2^{-3}, \dots, 2^{-m}$ (with the last m being that of Theorem 1), one finds a set of matrix units e_{ij} , $i, j = 1, \dots, 2^m$, which may be taken to approximately commute with any finite set in M_{n^∞} , such that $\tau\left(\sum_{i=1}^{2^m} e_{ii}\right) > 1 - \varepsilon$ and

$$\sigma(e_{ij}) \approx e_{i+1, j+1}$$

where the addition of indices is modulo 2^m (see [BEK] for details). Putting $e_i = e_{ii}$, Theorem 1 follows.

The proof of Corollary 2 from Theorem 1 now follows the lines of the proofs of [Voi, Lemma 3.3] or [BKRS, Lemma 2.1]. By [Voi, Lemma 3.1], for $D \in \mathcal{F}(M_{n^\infty})$ and $m \in \mathbb{N}$ there are $B_j \in \mathcal{F}(M_{n^\infty})$, $j = 0, 1, \dots, m$ with

$$\begin{aligned} B_0 &= B_m, \\ d(\sigma(B_i), B_{i+1}) &< \frac{5\pi}{m}, \quad i = 0, \dots, m-1, \\ D &\subseteq B_i, \quad i = 0, 1, \dots, m. \end{aligned}$$

If e_0, \dots, e_m is a Rohlin tower approximately commuting with B_0, \dots, B_m we may assume that $e = \sum_{k=0}^{m-1} e_k$ commute with D by a small perturbation of e , and putting

$$E \approx \sum_{i=0}^{m-1} B_i e_i,$$

E has the desired properties.

References

- [BEK] O. Bratteli, D. E. Evans and A. Kishimoto, The Rohlin property for quasifree automorphisms of the fermion algebra, under typing.
- [BKRS] O. Bratteli, A. Kishimoto, M. Rørdam and E. Størmer, The crossed product of a UHF algebra by a shift, *Preprint, July'92*.
- [CE] A. Connes and D. E. Evans, Embedding of $U(1)$ -current algebras in noncommutative algebras of classical statistical mechanics, *Commun. Math. Phys.* **12** (1989), 507–525.
- [Con] A. Connes, Outer conjugacy classes of automorphisms of factors, *Ann. Scient. Ec. Norm. Sup., 4 série*, **8** (1975), 383–420.
- [E] D. E. Evans, C^* -algebraic methods in statistical mechanics and field theory, *Inter. J. Modern Physics* **4** (1990), 1069–1118.
- [HO] R. H. Herman and A. Ocneanu, Stability for integer actions on UHF C^* -algebras, *J. Funct. Anal.* **59** (1984), 132–144.
- [Ren] J. Renault, A Groupoid Approach to C^* -algebras, *LNM 793*, Springer Verlag 1980.
- [Voi] D. Voiculescu, Almost inductive limit automorphisms and embeddings into AF-algebras, *Ergod. Th. & Dynam. Sys.* **6** (1986), 475–484.